

LIFTINGS OF GRADED QUASI-HOPF ALGEBRAS WITH RADICAL OF PRIME CODIMENSION

PAVEL ETINGOF AND SHLOMO GELAKI

1. INTRODUCTION

Let p be a prime, and let $RG(p)$ denote the set of equivalence classes of radically graded finite dimensional quasi-Hopf algebras over $\mathbb{C}$, whose radical has codimension p . In [EG1],[EG2] we completely describe the set $RG(p)$. Namely, we show that for $p > 2$, $RG(p)$ consists of the quasi-Hopf algebras $A(q)$ constructed in [G] for each primitive root of unity q of order p^2 , the Andruskiewitsch-Schneider Hopf algebras [AS1], and semisimple quasi-Hopf algebras $H_+(p)$ and $H_-(p)$ of dimension p ; on the other hand, $RG(2)$ consists of the Nichols Hopf algebras of dimension 2^n , $n \geq 1$, and the special quasi-Hopf algebras $H(2)$, $H_{\pm}(8) = A(\pm i)$, and $H(32)$ (of dimensions 2,8,32) constructed in [EG1].

The purpose of this paper is to classify finite dimensional quasi-Hopf algebras A whose radical is a quasi-Hopf ideal and has codimension p ; that is, A with $\text{gr}A \in RG(p)$, where $\text{gr}A$ is the associated graded algebra taken with respect to the radical filtration on A . The main result of this paper is the following theorem.

Theorem 1.1. *Let A be a finite dimensional quasi-Hopf algebra whose radical is a quasi-Hopf ideal of prime codimension p . Then either A is twist equivalent to a Hopf algebra, or it is twist equivalent to $H(2)$, $H_{\pm}(p)$, $A(q)$, or $H(32)$.*

Note that if A is a Hopf algebra (up to twist) then for $p = 2$ A is a Nichols Hopf algebra of dimension 2^n , while for $p > 2$ it is the dual to a finite dimensional pointed Hopf algebra with group of grouplike elements $\mathbb{Z}_p$. Such Hopf algebras are completely classified for $p \neq 5, 7$ in [AS2] (see Remark 1.10(v), [M]).

Note also that any finite tensor category whose simple objects are invertible and form a group of order p under tensor is the representation category of a quasi-Hopf algebra A as above. Thus this paper provides a classification of such categories.

The organization of the paper is as follows. Section 2 is devoted to preliminaries. In Section 3 we introduce a construction of semidirect products for quasi-Hopf algebras, which is used in the sequel. In Section 4 we consider liftings of the quasi-Hopf algebras $A(q)$ from [G], and of $H(32)$. We prove:

Theorem 1.2. *Let $\text{gr}(A)$ be $A(q)$ or $H(32)$. Then A is equivalent to $\text{gr}(A)$ as a quasi-Hopf algebra.*

In Section 5 we consider liftings of Andruskiewitsch-Schneider Hopf algebras [AS1]. We prove:

Theorem 1.3. *Let $\text{gr}(A)$ be an Andruskiewitsch-Schneider Hopf algebra. Then A is twist equivalent to a Hopf algebra.*

By [EG1], [EG2], Theorem 1.1 follows from Theorems 1.2 – 1.3.

Acknowledgments. The research of the first author was partially supported by the NSF grant DMS-9988796. The second author was supported by Technion V.P.R. Fund - Dent Charitable Trust - Non Military Research Fund, and by The Israel Science Foundation (grant No. 70/02-1). He also thanks MIT for its warm hospitality. Both authors were supported by BSF grant No. 2002040.

2. PRELIMINARIES

All constructions in this paper are done over the field of complex numbers $\mathbb{C}$.

We refer the reader to [D] for the definition of a quasi-Hopf algebra and a twist of a quasi-Hopf algebra.

2.1. The quasi-Hopf algebras $A(q)$.

Theorem 2.1. [G] *Let $n \geq 2$ be an integer.*

There exist n^3 -dimensional quasi-Hopf algebras $A(q)$, parametrized by primitive roots of unity q of order n^2 , which have the following structure. As algebras $A(q)$ are generated by a, x with the relations $ax = q^n xa$, $a^n = 1$, $x^{n^2} = 0$. The element a is grouplike, while the coproduct of x is given by the formula

$$\Delta(x) = x \otimes \sum_{y=0}^{n-1} q^y \mathbf{1}_y + 1 \otimes (1 - \mathbf{1}_0)x + a^{-1} \otimes \mathbf{1}_0 x,$$

where $\{\mathbf{1}_i | 0 \leq i \leq n-1\}$ is the set of primitive idempotents of $\mathbb{C}[a]$ defined by the condition $a\mathbf{1}_i = q^{ni}\mathbf{1}_i$. The associator is

$$\Phi := \sum_{i,j,k=0}^{n-1} q^{\frac{-i(j+k-(j+k)')}{n}} \mathbf{1}_i \otimes \mathbf{1}_j \otimes \mathbf{1}_k,$$

the distinguished elements are $\alpha = a$, $\beta = 1$, and the antipode is $S(a) = a^{-1}$, $S(x) = -x \sum_{z=0}^{n-1} q^{n-z} \mathbf{1}_z$.

Note that $S^2(x) = q^{n+1}x$, $S^2(a) = a$, so $S^{2n} = \text{Ad}(a)$.

2.2. The quasi-Hopf algebra $H(32)$.

Theorem 2.2. [EG1] *There exists a 32-dimensional quasi-Hopf algebra $H(32)$, which has the following structure. As an algebra $H(32)$ is generated by a, x, y with the relations $ax = -xa$, $ay = -ya$, $a^2 = 1$, $x^4 = 0$, $y^4 = 0$, $xy + iyx = 0$. The element a is grouplike, while the coproducts of x, y are given by the formulas*

$$\Delta(x) = x \otimes (p_+ + ip_-) + 1 \otimes p_+ x + a \otimes p_- x,$$

$$\Delta(y) = y \otimes (p_+ - ip_-) + 1 \otimes p_+ y + a \otimes p_- y,$$

where $p_+ := (1+a)/2$, $p_- := (1-a)/2$. The associator is $\Phi = 1 - 2p_- \otimes p_- \otimes p_-$, the distinguished elements are $\alpha = a$, $\beta = 1$, and the antipode is $S(a) = a$, $S(x) = -x(p_+ + ip_-)$, $S(y) = -y(p_+ - ip_-)$. Thus, $H(32)$ is generated by its quasi-Hopf subalgebras $A(i)$ and $A(-i)$ generated by a, x and a, y , respectively.

Note that $S^2(x) = ix$ and $S^2(y) = -iy$, so $S^4 = \text{Ad}(a)$.

2.3. Andruskiewitsch-Schneider Hopf algebras. In [AS1], Theorem 1.3, Andruskiewitsch and Schneider completely classified finite dimensional coradically graded Hopf algebras with coradical of prime dimension $p \geq 3$. Since this collection of Hopf algebras is obviously self-dual, the list of Theorem 1.3 of [AS1] also classifies radically graded Hopf algebras with radical of prime codimension $p \geq 3$.

Let us recall this list.

1. Quantum lines (Taft Hopf algebras) of dimension p^2 (type A_1). They are generated by a, x where

$$a^p = 1, x^p = 0, ax = qxa,$$

and

$$\Delta(a) = a \otimes a, \Delta(x) = x \otimes a + 1 \otimes x.$$

Here q is a primitive root of unity of degree p .

2. Quantum planes (book Hopf algebras) of dimension p^3 (type $A_1 \times A_1$). They are generated by a, x, y where

$$a^p = 1, x^p = 0, y^p = 0, xy - yx = 0, ax = qxa, ay = q^m ya$$

and

$$\Delta(a) = a \otimes a, \Delta(x) = x \otimes a + 1 \otimes x, \Delta(y) = y \otimes 1 + a^m \otimes y.$$

Here q is primitive a root of unity of degree p , and $m \in \mathbb{Z}_p$ is a nonzero element.

3. Algebras of type A_2, B_2, G_2 . They are generated by a, x, y satisfying the relations $axa^{-1} = qx$ and $aya^{-1} = q^b y$, where $b \in \mathbb{Z}_p$ is a nonzero element, and additional Serre relations coming from the Nichols algebra u_q^+ (of type A_2, B_2 , or G_2 , respectively) generated by x, y . The element a is grouplike, while the coproducts of x, y are given by the formulas

$$\Delta(x) = x \otimes a + 1 \otimes x, \Delta(y) = y \otimes a^d + 1 \otimes y,$$

where $0 \neq d \in \mathbb{Z}_p$, $b + d = -1$, and the following conditions hold:

- 1) The case A_2 : $p = 3$, or $p = 1$ modulo 3, and $d^2 + d + 1 = 0$.
- 2) The case B_2 : $p = 1$ modulo 4, and $2d^2 + 2d + 1 = 0$.
- 3) The case G_2 : $p = 1$ modulo 3, and $3d^2 + 3d + 1 = 0$.

Here, as before, q is a primitive root of unity of degree p .

4. Algebras of type $A_2 \times A_1$ ($p = 3$). They are generated by a, x, y as above (with $b = d = 1$), and a new element z such that $aza^{-1} = q^f z$,

$$\Delta(z) = z \otimes a^{-f} + 1 \otimes z,$$

where $f = 1$ or 2 (subject to appropriate Serre relations).

5. Algebras of type $A_2 \times A_2$ ($p = 3$). They are generated by a, x, y as above (with $b = d = 1$), and new elements x', y' , such that $ax'a^{-1} = q^f x'$, $ay'a^{-1} = q^f y'$,

$$\Delta(x') = x' \otimes a^{-f} + 1 \otimes x', \Delta(y') = y' \otimes a^{-f} + 1 \otimes y',$$

where $f = 1$ or 2 (subject to appropriate Serre relations).

In all cases the Hopf algebra is isomorphic $\mathbb{C}[\mathbb{Z}_p] \ltimes u_q^+$ as an algebra (where u_q^+ is the corresponding Nichols algebra).

2.4. Deformations. Let A be a finite dimensional quasi-Hopf algebra, in which the radical is a quasi-Hopf ideal. Let A_0 be the associated graded quasi-Hopf algebra of A under the radical filtration. The algebra A_0 has a grading $A_0 = \bigoplus_{i=0}^N A_0[i]$ by nonnegative integers. Thus we have gradings on the spaces of linear maps between tensor powers of A_0 . If f_0 is such a map of some degree d and f is obtained from f_0 by adding terms of degree higher than d , we will write $f = f_0 + \text{hdt}$ (where hdt stands for higher degree terms) and say that f is a deformation (or lifting) of f_0 .¹ Thus we can talk about deformations of algebra, coalgebra structures, associator, antipode, etc. In particular, we may say that A is a deformation of A_0 , in the sense that A is identified with A_0 as a graded vector space, so that the structure maps of A are given by some deformations of the structure maps of A_0 . This is the way we will think about A in the considerations below.

2.5. Quasi-Hopf liftings of Hopf algebras.

Proposition 2.3. *Let A be a finite dimensional quasi-Hopf algebra whose radical is a quasi-Hopf ideal, such that $A_0 = \text{gr}(A)$ is a Hopf algebra (i.e., the associator of A_0 is equal to 1). If $H^3(A_0^*, \mathbb{C}) = 0$ then A is twist equivalent to a Hopf algebra.*

Proof. Let Φ be the associator of A . We have $\Phi = 1 + \text{hdt}$. Write $\Phi = 1 + \gamma + \text{hdt}$, where γ is of degree d . Then γ is a 3-cocycle of A_0^* with coefficients in the trivial A_0^* -bimodule $\mathbb{C}$. Since $H^3(A_0^*, \mathbb{C}) = 0$, we have $\gamma = dj$, where $j \in A_0^{\otimes 2}[d]$, and the twisted associator Φ^{1+j} is equal to $1 +$ terms of degree $\geq d+1$. By continuing this procedure, we will come to a situation where $\Phi = 1$, as desired. $\square$

3. SEMIDIRECT PRODUCTS

Let H be a quasi-Hopf algebra with associator Φ , $g : H \rightarrow H$ an algebra automorphism. Define $\Delta_g(h) := (g \otimes g)(\Delta(g^{-1}(h)))$. Assume that

$$(1) \quad \Delta_g(h) = K \Delta(h) K^{-1}, \quad h \in H,$$

for some invertible element $K \in H \otimes H$ such that $(\varepsilon \otimes 1)(K) = (1 \otimes \varepsilon)(K) = 1$, and

$$g^{\otimes 3}(\Phi) = \Phi^K := (1 \otimes K)(id \otimes \Delta)(K)\Phi(\Delta \otimes id)(K^{-1})(K^{-1} \otimes 1).$$

Lemma 3.1. *The semidirect product algebra $\mathbb{C}[g, g^{-1}] \ltimes H$ is a quasi-Hopf algebra with associator Φ , $\Delta(g) = K^{-1}(g \otimes g)$ and $\varepsilon(g) = 1$.*

Proof. Using equation (1), we see that there exists a well defined coproduct on the algebra $\mathbb{C}[g, g^{-1}] \ltimes H$ which extends the coproduct on H and such that $\Delta(g) = K^{-1}(g \otimes g)$. It is straightforward to verify that $(\Delta \otimes id)\Delta(g) = \Phi^{-1}(id \otimes \Delta)\Delta(g)\Phi$. It is also easy to show that the antipode extends to the semidirect product. The lemma is proved. $\square$

Suppose further that $g^n(h) = aha^{-1}$ for some $a \in H$, and that

$$(2) \quad (g^{n-1})^{\otimes 2}(K) \dots g^{\otimes 2}(K)K = (a \otimes a)\Delta(a)^{-1}.$$

Lemma 3.2. *The ideal generated by $g^n - a$ is a quasi-Hopf ideal, so the quotient $\tilde{H} := (\mathbb{C}[g, g^{-1}] \ltimes H) / \langle g^n - a \rangle$ is a quasi-Hopf algebra.*

¹More generally, if $f_1, \dots, f_n$ have degrees $d_1, \dots, d_n$ then $f_1 + \dots + f_n + \text{hdt}$ will mean the sum of f_i plus a sum of corrections of degrees higher than all d_i .

Proof. Using condition (2) it is straightforward to see that

$$\begin{aligned} \Delta(g^n - a) &= \Delta(a)(a^{-1}g^n \otimes a^{-1}g^n - 1) \\ &= \Delta(a)(a^{-1}(g^n - a) \otimes a^{-1}g^n + 1 \otimes a^{-1}(g^n - a)). \end{aligned}$$

It is also straightforward to verify that $\langle g^n - a \rangle$ is invariant under the antipode. Thus $\langle g^n - a \rangle$ is a quasi-Hopf ideal. $\square$

4. PROOF OF THEOREM 1.2

Let $A_0 = A(q)$, where q is a primitive root of unity of order n^2 , or $A_0 = H(32)$ (in which case we set $n = 2$). Let A be a lifting of A_0 . We apply the semidirect product construction to the automorphism $g := S^2$ of the algebra A .

Proposition 1.2 of [D] provides a twist K such that $\Delta_g(y) = K\Delta(y)K^{-1}$, for all $y \in A$. Let K_0 be the degree zero part of K . Then K_0 commutes with $\Delta_0(y)$ for any $y \in A_0$, where Δ_0 is the comultiplication of A_0 . This implies, using straightforward computations, that $K_0 = 1$, and hence $K = 1 + \text{hdt}$.

Lemma 4.1. *The automorphism S^{2n} of A is inner.*

Proof. Let $G := \text{Aut}(A)/\text{Inn}(A)$ be the group of outer algebra automorphisms of A ; it is a linear algebraic group. Let S_0 be the antipode of A_0 . Then $S_0^{2n} = \text{Ad}(a)$. Hence the class $[S^{2n}]$ of S^{2n} is unipotent in G . On the other hand, from the main result of [ENO] (see also [HN]) we know that on the category of representations of A there exists an isomorphism of tensor functors $\theta_V : V \rightarrow \chi \otimes V^{****} \otimes \chi^{-1}$, where χ is a 1-dimensional representation of A . In our case, χ has order n . Therefore there exists a tensor isomorphism $c_V : V \rightarrow V^{**^{2n}}$, and thus S^{4n} is an inner automorphism of A : $S^{4n}(z) = czc^{-1}$ where $c \in A$ is the element realizing the automorphism c_V (i.e., $c_V = c|_V$). Hence, $[S^{2n}] = 1$ in G and S^{2n} is inner. $\square$

Thus, $S^{2n}(z) = bzb^{-1}$ for some element $b \in A$, where $b = a + \text{hdt}$ is a deformation of the grouplike element $a \in A_0$.

Lemma 4.2. *The element b can be chosen so that*

$$(3) \quad K^{-1}g^{\otimes 2}(K^{-1}) \dots (g^{n-1})^{\otimes 2}(K^{-1}) = \Delta(b)(b^{-1} \otimes b^{-1}).$$

Proof. Denote the left hand side of (3) by T .

Let c be the element defined above such that $S^{4n}(z) = czc^{-1}$. It is easy to check that $c = a^2 + \text{hdt}$.

We claim that

$$T(b \otimes b)T(b^{-1} \otimes b^{-1}) = \Delta(c)(c^{-1} \otimes c^{-1}).$$

Indeed, since $\{c_V\}$ is an isomorphism of tensor functors, $L(c_V \otimes c_W) = c_{V \otimes W}$, where L is the tensor structure on the functor $**^{2n}$. This defines an element $L \in A \otimes A$ such that $L = \Delta(c)(c^{-1} \otimes c^{-1})$. However, we know from [D] that the tensor structure on $**$ is given by K^{-1} , and hence the tensor structure on $**^{2n}$ is given by $K^{-1}(S^2 \otimes S^2)(K^{-1}) \dots (S^{4n-2} \otimes S^{4n-2})(K^{-1})$, which is equal to $T(S^{2n} \otimes S^{2n})(T) = T\text{Ad}(b \otimes b)(T)$, as desired.

Thus we have that $z := b^2c^{-1} = 1 + \text{hdt}$ is a central element. Replacing b with $bz^{-1/2}$, we may assume without loss of generality that b is chosen to satisfy the condition $z = 1$, i.e. $b^2 = c$.

We claim that for this choice of b , equation (3) holds. Indeed, assume the contrary, and let the lowest degree of nontrivial terms in the difference between the two sides of (3) be d . Let $\gamma \neq 0$ be the part of the difference that is homogeneous of degree d ; so $T = \Delta(b)(b^{-1} \otimes b^{-1}) - \gamma$ modulo elements of degree $\geq d+1$. Then modulo elements of degree $\geq d+1$, we have for any $y \in A$,

$$\begin{aligned} \Delta_{\text{Ad}(b)}(y) &= (b \otimes b)\Delta(b^{-1}yb)(b^{-1} \otimes b^{-1}) = T^{-1}\Delta(y)T \\ &= ((b \otimes b)\Delta(b)^{-1} + \gamma)\Delta(y)(\Delta(b)(b^{-1} \otimes b^{-1}) - \gamma). \end{aligned}$$

This means that for any $y \in A_0$ one has $[\gamma, \Delta_0(y)] = 0$, and in particular $[\gamma, a \otimes a] = 0$.

Also, modulo terms of degree $\geq d+1$,

$$(\Delta(b)(b^{-1} \otimes b^{-1}) - \gamma)(b \otimes b)(\Delta(b)(b^{-1} \otimes b^{-1}) - \gamma)(b^{-1} \otimes b^{-1}) = \Delta(c)(c^{-1} \otimes c^{-1}).$$

Since $b^2 = c$, this implies that

$$\gamma + (a \otimes a)\gamma(a^{-1} \otimes a^{-1}) = 0.$$

Since $[\gamma, a \otimes a] = 0$, this is equivalent to $2\gamma = 0$. Contradiction. $\square$

Lemma 3.2 and Lemma 4.2 imply that we can define the semidirect product quasi-Hopf algebra

$$\tilde{H} := (\mathbb{C}[g, g^{-1}] \ltimes A) / \langle g^n - b \rangle;$$

it is of dimension n^4 for $A = A(q)$, and 64 for $A = H(32)$.

Let $J \in \tilde{H} \otimes \tilde{H}$ be a deformation of the twist J_0 from $[G]$. Then it follows from $[G]$ that $H := \tilde{H}^{J^{-1}}$ is a quasi-Hopf lifting (=deformation) of a Hopf algebra H_0 . If $A_0 = A(q)$ then H_0 is the Taft Hopf algebra of dimension n^4 . On the other hand, if $A_0 = H(32)$ then H_0 is a book Hopf algebra of dimension 64, generated by g, x_+, x_- satisfying $gx_+g^{-1} = ix_+$, $gx_-g^{-1} = -ix_-$, $x_+x_- = x_-x_+$. The element g is grouplike and

$$\Delta(x_+) = x_+ \otimes g + 1 \otimes x_+, \quad \Delta(x_-) = x_- \otimes 1 + g^{-1} \otimes x_-.$$

Theorem 4.3. *If H is a basic quasi-Hopf algebra such that $H_0 = grH$ then there exists a quasi-Hopf twist F such that $H \cong H_0^F$ and $F = 1$ modulo higher terms.*

Proof. Let us first show that H is twist equivalent to a Hopf algebra. By Proposition 2.3, for this it is sufficient to show that $H^3(H_0^*, \mathbb{C}) = 0$. If H_0 is the Taft algebra then $H_0^* \cong H_0$ and one finds (see [GK]), that $H^\bullet(H_0^*, \mathbb{C})$ is the polynomial algebra on an element of degree 2. In particular, $H^3(H_0, \mathbb{C}) = 0$, as required. On the other hand, if H_0 is the book algebra of dimension 64 then H_0^* is generated by g, x, y with $g^4 = 1$, $x^4 = y^4 = xy - yx = 0$, $gxg^{-1} = ix$, $gyg^{-1} = iy$. From this it is easy to obtain using Künneth formula that $H^\bullet(H_0^*, \mathbb{C})$ is the polynomial algebra in two elements of degree 2 (see [GK]). In particular, again $H^3(H_0^*, \mathbb{C}) = 0$.

Now, it is easy to see, using the methods of [AS2], that in both cases (the Taft algebra and the book algebra), H_0 has no non-trivial Hopf liftings, so there exists a twist $F = 1 + \text{hdt}$ and an isomorphism of quasi-Hopf algebras $\eta : H_0^F \rightarrow H$, as desired. ² $\square$

²We note that if H_0 is the book algebra of dimension 64, then H_0^* , unlike H_0 , has a nontrivial Hopf lifting, which is the dual of the quantum sl_2 at the 4-th root of unity.

Replacing J with JF , we may assume that $F = 1$. Thus we find that $\tilde{H} = H_0^J$. So we have a homomorphism of quasi-Hopf algebras

$$(4) \quad \eta : A \rightarrow H_0^J.$$

The map η may be regarded as an injective linear map $A_0 \rightarrow H_0^J$, whose image is a quasi-Hopf subalgebra of H_0^J , and whose degree 0 part is the “tautological” homomorphism $\eta_0 : A_0 \rightarrow H_0^{J_0^3}$. Let E be the set of pairs (η, J) satisfying this condition. On E there is an action of the “gauge group” $\mathcal{G}$, generated by the following three types of transformations.

1. $(\eta, J) \rightarrow (\eta \circ \xi, J)$, where $\xi = 1 + \text{hdt}$ is a linear automorphism of A_0 .
2. $(\eta, J) \rightarrow (\text{Ad}(h) \circ \eta, (h \otimes h)J\Delta(h)^{-1})$, where $h = 1 + \text{hdt}$ is an element of H_0 , and Δ is the coproduct of H_0 .
3. $(\eta, J) \rightarrow (\eta, TJ)$, where $T = 1 + \text{hdt}$ belongs to $\eta(A_0)^{\otimes 2}$.

It is clear that the transformations from $\mathcal{G}$ do not change the equivalence class of the quasi-Hopf algebra $\eta(A_0)$. Thus, Theorem 1.2 follows from the following proposition.

Proposition 4.4. *Any pair $(\eta, J) \in E$ is equivalent under $\mathcal{G}$ to (η_0, J_0) .*

Proof. It suffices to prove that for any $d > 0$, any pair $(\eta, J) \in E$ can be brought by the action of $\mathcal{G}$ to a pair equal to (η_0, J_0) modulo terms of degree $\geq d$. We will prove it by induction in d . The base of induction is obvious. To prove the induction step, we need to take a pair (η, J) which is equal to (η_0, J_0) modulo terms of degree d , and show it is equivalent to a pair equal to (η_0, J_0) modulo terms of degree $d+1$.

Thus, the multiplication in A_0 obtained by pullback of the multiplication of H_0 by η is given by $z * w = zw + c_d(z, w) + \text{hdt}$, where zw is the multiplication in A_0 , and c_d is a bilinear operator $A_0 \times A_0 \rightarrow A_0$ of degree d , which is a Hochschild 2-cocycle. Similarly, $\eta(z) = \eta_0(z) + \eta_d(z) + \text{hdt}$, where η_d is a degree d operator $A_0 \rightarrow H_0$, and $J = TJ_0$, $T = 1 + t_d + \text{hdt}$, where $t_d \in H_0 \otimes H_0$ has degree d .

We have $\eta(z * w) = \eta(z)\eta(w)$ which is equivalent to

$$(\eta_0 + \eta_d + \cdots)(zw + c_d(z, w) + \cdots) = (\eta_0 + \eta_d + \cdots)(z)(\eta_0 + \eta_d + \cdots)(w).$$

Taking z of degree d_z and w of degree d_w , and looking at terms of degree $d + d_z + d_w$ we get $\eta_d(zw) + \eta_0(c_d(z, w)) = \eta_d(z)\eta_0(w) + \eta_0(z)\eta_d(w)$, which implies $\eta_0(c_d(z, w)) = \eta_d(z)\eta_0(w) + \eta_0(z)\eta_d(w) - \eta_d(zw)$. Therefore, $\eta_0(c_d)$ is the differential of a 1-cochain $\eta_d \in C^1(A_0, H_0)$. So η_0 sends the cohomology class of $c_d \in H^2(A_0, A_0)$ into 0 in $H^2(A_0, H_0)$. Since A_0 is a direct summand in H_0 as an A_0 -bimodule, this implies that the class $[c_d] \in H^2(A_0, A_0)$ is zero. Thus, c_d is the differential of some element τ of degree d in $C^1(A_0, A_0)$. So, using an element $g_1 \in \mathcal{G}$ of type 1, we can replace (η, J) with an equivalent pair for which $c_d = 0$. Thus in the sequel we may assume that $c_d = 0$.

Now, η_d is a Hochschild 1-cocycle of A_0 with coefficients in H_0 . Consider the cohomology class $[\eta_d]$ in $H^1(A_0, H_0)$. We claim that in fact, $[\eta_d]$ belongs to the subspace $H^1(A_0, A_0) \subset H^1(A_0, H_0)$. To prove this, note that since $\eta(A_0)$ is closed under coproduct, for any $z \in A_0$ we have

$$\Delta_0(\eta_d(z)) - (\eta_d \otimes \eta_0 + \eta_0 \otimes \eta_d)(\Delta_0(z)) = [\Delta_0(z), t] \text{ mod } A_0 \otimes A_0,$$

where Δ_0 denotes the coproduct of $H_0^{J_0}$ and A_0 , and $t := t_d$.

³Note the difference between the map $\eta_0 : A_0 \rightarrow H_0$ and the map $A_0 \rightarrow H_0^*$ (not $H_0!$) constructed in [EG1].

Now recall that $H_0 = \bigoplus_{k=0}^{n-1} A_0 g^k$. Thus, $\eta_d = \sum_{k=0}^{n-1} \omega_k$, where $\omega_k \in Z^1(A_0, A_0 g^k)$ (a 1-cocycle). Similarly, $t = \sum_{k,l=0}^{n-1} t_{kl}$, $t_{kl} \in A_0 g^k \otimes A_0 g^l$. Thus for $k \neq 0$ we have

$$-(\omega_k \otimes \eta_0)(\Delta_0(z)) = [\Delta_0(z), t_{k0}]$$

Applying the counit in the second component, we get that $\omega_k(z) = [z, y_k]$ for some $y_k \in A_0 g^k$, which implies that ω_k is an inner derivation.

Thus $[\eta_d] \in H^1(A_0, A_0)$, and hence we could have chosen g_1 so that $[\eta_d] = 0$. So we may assume $[\eta_d] = 0$ and thus η_d is the bracketing operator with an element $y \in H_0$. Thus using an appropriate element $g_2 \in \mathcal{G}$ of type 2, we can assume in the sequel that $\eta_d = 0$.

Now we will show that in the situation when $c_d = 0$ and $\eta_d = 0$, we must have $t \in A_0 \otimes A_0$. Then by using an element $g_3 \in \mathcal{G}$ of type 3, we can come to a situation with $t = 0$ which completes the induction step.

Since the associator $\Phi := \Phi_0^T$ lies in $\eta(A_0)^{\otimes 3}$, we find that

$$(5) \quad t \otimes 1 + (\Delta_0 \otimes id)(t) - \Phi_0(id \otimes \Delta_0)(t)\Phi_0^{-1} + \Phi_0(1 \otimes t)\Phi_0^{-1} \in A_0^{\otimes 3}.$$

Let $H_0 = A_0 \oplus H'$, where H' is the direct sum of $A_0 g^k$ for $k = 1, \dots, n-1$. Let t' be the projection of t to $H' \otimes H_0$. Since the first component of $t - t'$ is in A_0 , we get from (5):

$$(6) \quad t' \otimes 1 + (\Delta_0 \otimes id)(t') - \Phi_0(id \otimes \Delta_0)(t')\Phi_0^{-1} + \Phi_0(1 \otimes t')\Phi_0^{-1} \in A_0 \otimes H_0 \otimes H_0.$$

But out of the four summands in this expression, all but the last one belong to $H' \otimes H_0 \otimes H_0$, while the last summand is in $A_0 \otimes H_0 \otimes H_0$. Hence the last summand is zero and $t' = 0$. Similarly, one shows that the projection t'' of t to $H_0 \otimes H'$ is zero. Hence, $t \in A_0 \otimes A_0$, as desired. $\square$

The theorem is proved.

5. PROOF OF THEOREM 1.3

Let H be a Hopf algebra from the list of Subsection 2.3. Since this list is self-dual, Theorem 1.3 follows from Proposition 2.3 and the following proposition.

Proposition 5.1. *We have $H^3(H, \mathbb{C}) = 0$.*

Proof. By a standard argument, $H^3(H, \mathbb{C}) = H^3(\mathbb{C}[\mathbb{Z}_p] \ltimes \mathfrak{u}_q^+, \mathbb{C}) = H^3(\mathfrak{u}_q^+, \mathbb{C})^{\mathbb{Z}_p}$.

Let us first assume that $p > 3$. In this case, according to Section 2.3, the rank of the corresponding simple Lie algebra is ≤ 2 , and p is greater than its Coxeter number. Thus we can apply [GK], Theorem 2.5. In this theorem, it is shown that $H^\bullet(\mathfrak{u}_q^+, \mathbb{C}) = \bigoplus_{w \in W} \mathbb{C} \eta_w \otimes S(\mathfrak{n}_+)$, where $S(\mathfrak{n}_+)$ is the symmetric algebra of the positive part of the associated Lie algebra, sitting in degree 2, W is the Weyl group, and η_w has degree $l(w)$ (=length of w).

The action of the generator a of $\mathbb{Z}_p$ on $H^\bullet(\mathfrak{u}_q^+, \mathbb{C})$ is given as follows: a acts trivially on $S(\mathfrak{n}_+)$ and $a \circ \eta_w = \lambda_w \eta_w$, where in the A_1 -case $\lambda_w = q^{-1}$ for the nontrivial element $w \in W$, and in the other (rank 2) cases $\lambda_w = q^{-(m+nd)}$, where m, n are determined by writing the weight $\gamma_w := \sum_{\alpha \in R^+, w(\alpha) < 0} \alpha = \rho - w(\rho)$ as an integral combination $m\alpha_1 + n\alpha_2$ of the simple roots (here ρ is half the sum of positive roots, and R_+ is the set of positive roots; in the non-simply laced case, α_2 always denotes the short root).

We claim that $H^3(\mathfrak{u}_q^+, \mathbb{C})^{\mathbb{Z}_p} = 0$. Indeed, since $\mathbb{Z}_p$ acts trivially on $S(\mathfrak{n}_+)$, and since $\lambda_w \neq 1$ if w is a simple reflection, we have $H^3(\mathfrak{u}_q^+, \mathbb{C})^{\mathbb{Z}_p} = (\bigoplus_{w \in W, l(w)=3} \mathbb{C} \eta_w)^{\mathbb{Z}_p}$.

We will now show that in each of the four cases $A_1 \times A_1$, A_2 , B_2 and G_2 , $\lambda_w \neq 1$, and hence $H^3(\mathfrak{u}_q^+, \mathbb{C})^{\mathbb{Z}_p} = 0$.

In the case $A_1 \times A_1$ there are no elements w of length 3, so the result is clear.

In the case A_2 , there is one element w of length 3 and one computes that $\gamma_w = 2\alpha_1 + 2\alpha_2$. Therefore $\lambda_w = (q^{-1})^{2+2d}$. Since $d^2 + d + 1 = 0$, $d + 1 \neq 0$, so $\lambda_w \neq 1$.

In the case B_2 , there are two elements w_1, w_2 of length 3 and one computes that $\gamma_{w_1} = 3\alpha_1 + 3\alpha_2$, $\gamma_{w_2} = 2\alpha_1 + 4\alpha_2$. Therefore $\lambda_{w_1} = (q^{-1})^{3+3d}$ and $\lambda_{w_2} = (q^{-1})^{2+4d}$. Since $2d^2 + 2d + 1 = 0$, we have $d + 1, 2d + 1 \neq 0$, so $\lambda_{w_1}, \lambda_{w_2} \neq 1$.

In the case G_2 , there are two elements w_1, w_2 of length 3 and one computes that $\gamma_{w_1} = 4\alpha_1 + 4\alpha_2$, $\gamma_{w_2} = 2\alpha_1 + 6\alpha_2$. Therefore $\lambda_{w_1} = (q^{-1})^{4+4d}$ and $\lambda_{w_2} = (q^{-1})^{2+6d}$. Since $3d^2 + 3d + 1 = 0$, we have $d + 1, 3d + 1 \neq 0$, so $\lambda_{w_1}, \lambda_{w_2} \neq 1$. Thus the result for $p > 3$ follows.

Finally, let us consider the case $p = 3$. In the rank 1 case the above argument applies, so we consider rank ≥ 2 . Then the argument above does not quite apply, since the rank of the Lie algebra can now be bigger than 2, and p may be equal to the Coxeter number (a case not covered by Theorem 2.5 in [GK]). Thus we will use a slightly different argument. By [GK], the Nichols algebra $\mathfrak{u}_q^+$ has a filtration under which $H^\bullet(\text{gru}_q^+, \mathbb{C}) = \Lambda_q \otimes S(\mathfrak{n}_+)$, where Λ_q is the q -analogue of the exterior algebra generated by ε_α , α a positive root, such that $\varepsilon_\alpha \varepsilon_\beta + q^{<\alpha, \beta>} \varepsilon_\beta \varepsilon_\alpha = 0$ for $\beta < \alpha$ and $\varepsilon_\alpha^2 = 0$ for any α .

The generator a of $\mathbb{Z}_p$ acts trivially on $S(\mathfrak{n}_+)$ and by $a\varepsilon_\alpha a^{-1} = q^{-(m+nd)}\varepsilon_\alpha$, where $\alpha = m\alpha_1 + n\alpha_2$. The group G of grouplikes in the Frobenius-Lusztig quantum group acts on both $H^\bullet(\text{gru}_q^+, \mathbb{C})$ and $H^\bullet(\mathfrak{u}_q^+, \mathbb{C})$, and the second one is isomorphic to a subrepresentation of the first one by a spectral sequence argument [GK].

In the case A_2 ($d = 1$) we claim that $H^3(\text{gru}_q^+, \mathbb{C})^{\mathbb{Z}_3} = 0$. Indeed, we have $H^3(\text{gru}_q^+, \mathbb{C}) = \langle \varepsilon_{\alpha_1} \varepsilon_{\alpha_2} \varepsilon_{\alpha_1 + \alpha_2} \rangle$, and the generator a of $\mathbb{Z}_p$ acts on this 1-dimensional space by $q^{-(1+d+1+d)} = q^{-4} \neq 1$.

In the case $A_2 \times A_1$ ($d = 1$), we have that the Nichols algebra $\mathfrak{u}_q^+$ is generated by x, y, z with $aza^{-1} = q^f z$. Thus, if β is the simple root corresponding to z , we have, $H^3(\text{gru}_q^+, \mathbb{C})^{\mathbb{Z}_3} = \langle \varepsilon_\beta \varepsilon_{\alpha_1} \varepsilon_{\alpha_2} \rangle$ if $f = 2$, and is zero if $f = 1$.

In the case $A_2 \times A_2$ ($d = 1$), we have that the Nichols algebra $\mathfrak{u}_q^+$ is generated by x, y, x', y' with $axa^{-1} = qx$, $aya^{-1} = qy$, $ax'a^{-1} = q^f x'$ and $ay'a^{-1} = q^f y'$. Hence, if β_1, β_2 are the roots attached to x', y' , we find that

$$H^3(\text{gru}_q^+, \mathbb{C})^{\mathbb{Z}_3} = \langle \varepsilon_{\beta_1} \varepsilon_{\alpha_1} \varepsilon_{\alpha_2}, \varepsilon_{\alpha_1} \varepsilon_{\beta_1} \varepsilon_{\beta_2} \mid i = 1, 2 \rangle$$

if $f = 1$, and is zero if $f = 2$.

But even in cases where $H^3(\text{gru}_q^+, \mathbb{C})^{\mathbb{Z}_3} \neq 0$, this cohomology is killed in the spectral sequence $H^\bullet(\text{gru}_q^+, \mathbb{C})^{\mathbb{Z}_3} \rightarrow H^\bullet(\mathfrak{u}_q^+, \mathbb{C})^{\mathbb{Z}_3}$ since by [GK], the only weights of terms under G which can survive in this sequence must be of the form $w(\rho) - \rho$ while $-(\alpha_1 + \alpha_2) \neq w(\rho) - \rho$ modulo 3. So in all cases $H^3(\mathfrak{u}_q^+, \mathbb{C})^{\mathbb{Z}_3} = 0$, and the result follows. $\square$

REFERENCES

- [AS1] N. Andruskiewitsch and H-J. Schneider, Finite quantum groups and Cartan matrices, *Adv. Math.* **154** (2000), no. 1, 1–45.
- [AS2] N. Andruskiewitsch and H-J. Schneider, Finite quantum groups over abelian groups of prime exponent, *Ann. Sci. Ecole Norm. Sup. (4)* **35** (2002), no. 1, 1–26.
- [D] V. Drinfeld, Quasi-Hopf algebras. (Russian) *Algebra i Analiz* **1** (1989), no. 6, 114–148; translation in *Leningrad Math. J.* **1** (1990), no. 6, 1419–1457.

- [EG1] P. Etingof and S. Gelaki, Finite-dimensional quasi-Hopf algebras with radical of codimension 2, *Mathematical Research Letters* **11** (2004), 685–696.
- [EG2] P. Etingof and S. Gelaki, On radically graded finite dimensional quasi-Hopf algebras, submitted, math.QA/0403096.
- [ENO] P. Etingof, D. Nikshych and V. Ostrik, An analogue of Radford’s S^4 formula for finite tensor categories, *Int. Math. Res. Not.*, to appear, math.QA/0404504.
- [G] S. Gelaki, Basic quasi-Hopf algebras of dimension n^3 , *Journal of Pure and Applied Algebra*, to appear, math.QA/0402159.
- [GK] V. Ginzburg and S. Kumar, Cohomology of quantum groups at roots of unity, *Duke Math. J.* **69** (1993), no.1, 179–198.
- [HN] F. Hausser and F. Nill, Integral Theory for Quasi-Hopf Algebras, math.QA/9904164.
- [M] I. M. Musson, Finite quantum groups and pointed Hopf algebras. Quantum groups and Lie theory (Durham, 1999), 191–205, *London Math. Soc. Lecture Note Ser.*, **290**, Cambridge Univ. Press, Cambridge, 2001.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

E-mail address: `etingof@math.mit.edu`

DEPARTMENT OF MATHEMATICS, TECHNION-ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 32000, ISRAEL

E-mail address: `gelaki@math.technion.ac.il`